



# The Australian Journal of Mathematical Analysis and Applications

<http://ajmaa.org>

Volume 2, Issue 1, Article 11, pp. 1-10, 2005



---

## ON THE ULAM STABILITY FOR EULER-LAGRANGE TYPE QUADRATIC FUNCTIONAL EQUATIONS

MATINA JOHN RASSIAS AND JOHN MICHAEL RASSIAS

*Received 16 November, 2004; accepted 2 February, 2004; published 30 May, 2005.*

STATISTICS AND MODELLING SCIENCE, UNIVERSITY OF STRATHCLYDE, LIVINGSTONE TOWER, 26  
RICHMOND STR, GLASGOW, UK, G1 1XH

PEDAGOGICAL DEPARTMENT, E. E., NATIONAL AND CAPODISTRIAN UNIVERSITY OF ATHENS, SECTION OF  
MATHEMATICS AND INFORMATICS, 4, AGAMEMNONOS STR, AGHIA PARASKEVI, ATHENS 15342, GREECE

**ABSTRACT.** In 1940 (and 1968) S. M. Ulam proposed the well-known Ulam stability problem. In 1941 D.H. Hyers solved the Hyers-Ulam problem for linear mappings. In 1951 D. G. Bourgin has been the second author treating the Ulam problem for additive mappings. In 1978 according to P.M. Gruber this kind of stability problems is of particular interest in probability theory and in the case of functional equations of different types. In 1982-2004 we established the Hyers-Ulam stability for the Ulam problem for different mappings. In 1992-2000 J.M. Rassias investigated the Ulam stability for Euler-Lagrange mappings. In this article we solve the Ulam problem for Euler-Lagrange type quadratic functional equations. These stability results can be applied in mathematical statistics, stochastic analysis, algebra, geometry, as well as in psychology and sociology.

*Key words and phrases:* Ulam stability, Euler-Lagrange type mapping, Quadratic equation.

2000 *Mathematics Subject Classification.* Primary 39B. Secondary 26D.

---

ISSN (electronic): 1449-5910

© 2005 Austral Internet Publishing. All rights reserved.

## 1. INTRODUCTION

In 1940 (and 1968) S. M. Ulam [28] proposed *the Ulam stability problem*:

“When is it true that by slightly changing the hypotheses of a theorem one can still assert that the thesis of the theorem remains true or approximately true ?”

In particular he stated *the stability question*:

“Let  $G_1$  be a group and  $G_2$  a metric group with the metric  $\rho(\cdot, \cdot)$ . Given a constant  $\delta > 0$ , does there exist a constant  $c > 0$  such that if a mapping  $f : G_1 \rightarrow G_2$  satisfies  $\rho(f(xy), f(x)f(y)) < c$  for all  $x, y \in G_1$ , then a unique homomorphism  $h : G_1 \rightarrow G_2$  exists with  $\rho(f(x), h(x)) < \delta$  for all  $x \in G_1$ ?”

In 1941 D.H. Hyers [13] solved this problem for linear mappings. In 1951, D.G. Bourgin [3] was the second author to treat the Ulam problem for additive mappings. In 1978, according to P.M. Gruber [12], this kind of stability problems is of particular interest in probability theory and in the case of functional equations of different types. In 1980 and in 1987, I. Fenyő [7], [8] established the stability of the Ulam problem for quadratic and other mappings. In 1987 Z. Gajda and R. Ger [10] showed that one can get analogous stability results for subadditive multifunctions. Other interesting stability results have been achieved also by the following authors: J. Aczél [1], C. Borelli and G.L. Forti [2], [9], P.W. Cholewa [4], St. Czerwik [5], and H. Drljevic [6], and Pl. Kannappan [15]. In 1982-2004, J.M. Rassias [16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27] solved the above Ulam problem for different mappings. In 1999, P. Găvruta [11] answered a question of J.M. Rassias [18] concerning the stability of the Cauchy equation. In 1998, S.-M. Jung [14] and in 2002-2003 we [25, 26] investigated the Hyers-Ulam stability for additive and quadratic mappings on restricted domains. In 1992-2000 the second author ([19, 21, 22, 23, 24]) investigated the Ulam stability for Euler-Lagrange mappings.

In this article we solve the Ulam stability problem for Euler-Lagrange type quadratic functional equations.

Throughout this paper, let  $X$  be a real normed space and  $Y$  be a real Banach space in the case of functional inequalities, as well as let  $X$  and  $Y$  be real linear spaces for functional equations.

Let us introduce the Euler-Lagrange type quadratic functional equation

$$(1.1) \quad \begin{aligned} & Q(m_1 a_1 x_1 + m_2 a_2 x_2) + m_1 m_2 Q(a_2 x_1 - a_1 x_2) \\ &= (m_1 a_1^2 + m_2 a_2^2) [m_1 Q(x_1) + m_2 Q(x_2)] \end{aligned}$$

with mappings  $Q : X \rightarrow Y$ , for all  $x_1, x_2 \in X$ , and any fixed pair  $(a_1, a_2)$  of reals  $a_i \neq 0$  and any fixed pair  $(m_1, m_2)$  of positive reals  $m_i (i = 1, 2) : 0 < m = \frac{m_1 + m_2}{m_1 m_2 + 1} (m_1 a_1^2 + m_2 a_2^2) \neq 1$ .

**Definition 1.1.** A mapping  $Q : X \rightarrow Y$  is called *Euler-Lagrange type quadratic*, if the above-mentioned functional equation (1.1) holds for every  $(x_1, x_2, \dots, x_p) \in X^p$  with an arbitrary but fixed  $p = 2, 3, 4, \dots$

In this paper, we establish an approximation of approximately Euler-Lagrange type quadratic mappings  $f : X \rightarrow Y$  by Euler-Lagrange type quadratic mappings  $Q : X \rightarrow Y$ , such that the corresponding functional inequality

$$(1.2) \quad \begin{aligned} & \|f(m_1 a_1 x_1 + m_2 a_2 x_2) + m_1 m_2 f(a_2 x_1 - a_1 x_2) \\ & - (m_1 a_1^2 + m_2 a_2^2) [m_1 f(x_1) + m_2 f(x_2)]\| \\ & \leq c \end{aligned}$$

holds with a constant  $c \geq 0$  (independent of  $x_1, x_2 \in X$ ).

It is useful for the following, to observe that, from (1.1) with  $x_1 = x_2 = 0$ , and  $0 < m \neq 1$ , we get  $(m_1 m_2 + 1) |1 - m| Q(0) = 0$ , or

$$(1.3) \quad Q(0) = 0.$$

Similarly, from (1.2), one finds  $(m_1 m_2 + 1) |1 - m| \|f(0)\| \leq c$ , or

$$(1.4) \quad \|f(0)\| \leq \frac{c}{(m_1 m_2 + 1) |1 - m|} = \frac{c}{m_1 m_2 + 1} \begin{cases} \frac{1}{m-1}, & \text{if } m > 1 \\ \frac{1}{1-m}, & \text{if } 0 < m < 1. \end{cases}$$

Let us denote

$$(1.5) \quad \bar{Q}(x) = m_0 \begin{cases} \frac{m_1 Q\left(\frac{a_1}{m_0}x\right) + m_2 Q\left(\frac{a_2}{m_0}x\right)}{m}, & \text{if } m > 1 \\ m \left[ m_1 Q\left(\frac{b_1}{m_0}x\right) + m_2 Q\left(\frac{b_2}{m_0}x\right) \right], & \text{if } 0 < m < 1, \end{cases}$$

and  $b_i = \frac{a_i}{m}$  ( $i = 1, 2$ ) and  $m_0 = \frac{(m_1 + m_2)}{(m_1 m_2 + 1)}$ , as well as

$$(1.6) \quad \bar{\bar{Q}}(x) = \frac{1}{m_0 m_1} \begin{cases} \frac{Q(m_1 a_1 x) + m_1 m_2 Q(a_2 x)}{m}, & \text{if } m > 1 \\ m [Q(m_1 b_1 x) + m_1 m_2 Q(b_2 x)], & \text{if } 0 < m < 1, \end{cases}$$

for all  $x \in X$ .

**Definition 1.2.** Let  $X$  and  $Y$  be real linear spaces and  $m > 1$ . Then the following equation

$$(1.7) \quad F^a(Q) = Q(m_1 a_1 x) + m_1 m_2 Q(a_2 x) - m_0^2 m_1 \left[ m_1 Q\left(\frac{a_1}{m_0}x\right) + m_2 Q\left(\frac{a_2}{m_0}x\right) \right] = 0,$$

is called *fundamental functional equation of first type*. This (1.7) is equivalent to

$$(1.8) \quad (M^a(Q) =) \frac{F^a(Q)}{m_0 m_1 m} = \bar{\bar{Q}}(x) - \bar{Q}(x) = 0, m > 1.$$

Note that if  $X$  and  $Y$  are real normed linear spaces and  $m > 1$ , then

$$(1.9) \quad \|F^a(f)\| \leq \varepsilon_1,$$

with a constant  $\varepsilon_1 \geq 0$  (independent of  $x_1, x_2 \in X$ ). This inequality (1.9) is equivalent to

$$(1.10) \quad (\|M_a(f)\| =) \frac{1}{m_0 m_1 m} \|F_a(f)\| = \|\bar{\bar{f}}(x) - \bar{f}(x)\| \leq \frac{\varepsilon_1}{m_0 m_1 m}, m > 1.$$

**Definition 1.3.** Let  $X$  and  $Y$  be real linear spaces,  $b_i = \frac{a_i}{m}$  ( $i = 1, 2$ ) and  $0 < m < 1$ . Then

$$(1.11) \quad F^b(Q) = Q(m_1 b_1 x) + m_1 m_2 Q(b_2 x) - m_0^2 m_1 \left[ m_1 Q\left(\frac{b_1}{m_0}x\right) + m_2 Q\left(\frac{b_2}{m_0}x\right) \right] = 0$$

is called *fundamental functional equation of second type*. This (1.11) is equivalent to

$$(1.12) \quad (M^b(Q) =) \frac{m}{m_0 m_1} F^b(Q) = \bar{\bar{Q}}(x) - \bar{Q}(x) = 0, 0 < m < 1.$$

Note that if  $X$  and  $Y$  are real normed linear spaces and  $0 < m < 1$ , then

$$(1.13) \quad \|F^b(f)\| \leq \varepsilon_2,$$

with a constant  $\varepsilon_2 \geq 0$  (independent of  $x_1, x_2 \in X$ ). This inequality (1.13) is equivalent to

$$(1.14) \quad (\|M^b(f)\| =) \frac{m}{m_0 m_1} \|F^b(f)\| = \|\bar{f}(x) - \bar{f}(x)\| \leq \frac{m}{m_0 m_1} \varepsilon_2, \quad 0 < m < 1.$$

Now, claim that for  $n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\} = \{0, 1, 2, \dots\}$

$$(1.15) \quad Q(x) = \begin{cases} m^{-2n} Q(m^n x), & \text{if } m > 1 \\ m^{2n} Q(m^{-n} x), & \text{if } 0 < m < 1 \end{cases}$$

holds for all  $x \in X$ .

Let us consider first the case  $m \geq 1$ . For  $n = 0$ , it is trivial. From (1.3), (1.5) and (1.1), with  $x_i = \frac{a_i}{m_0} x$  ( $i = 1, 2$ ), we obtain

$$Q(mx) = mm_0 \left[ m_1 Q\left(\frac{a_1}{m_0} x\right) + m_2 Q\left(\frac{a_2}{m_0} x\right) \right],$$

or

$$(1.16) \quad \bar{Q}(x) = m^{-2} Q(mx).$$

Besides from (1.3), (1.6) and (1.1), with  $x_1 = x, x_2 = 0$ , one gets

$$Q(m_1 a_1 x) + m_1 m_2 Q(a_2 x) = mm_0 m_1 Q(x),$$

or

$$(1.17) \quad \bar{\bar{Q}}(x) = Q(x).$$

Therefore from (1.8), (1.16) and (1.17) we have

$$(1.18) \quad Q(x) = m^{-2} Q(mx),$$

which is (1.15) for  $n = 1$ , and  $m > 1$ . Assume (1.15),  $m > 1$ , is true and from (1.18), with  $m^n x$  on place of  $x$ , we get :

$$(1.19) \quad Q(m^{n+1}x) = m^2 Q(m^n x) = m^2 (m^n)^2 Q(x) = (m^{n+1})^2 Q(x).$$

This formula (1.19), by induction, proves formula (1.15) for  $m > 1$ .

Let us consider now the case  $0 < m < 1$ . Similarly from (1.3), (1.5) and (1.1), with  $x_i = \frac{a_i}{m_0} \frac{x}{m}$  ( $i = 1, 2$ ), we obtain

$$Q(x) = mm_0 \left[ m_1 Q\left(\frac{a_1}{m_0} \frac{x}{m}\right) + m_2 Q\left(\frac{a_2}{m_0} \frac{x}{m}\right) \right],$$

or

$$(1.20) \quad Q(x) = \bar{Q}(x).$$

Besides from (1.3), (1.6) and (1.1), with  $x_1 = \frac{x}{m}, x_2 = 0$ , one gets

$$Q\left(\frac{m_1 a_1}{m} x\right) + m_1 m_2 Q\left(\frac{a_2}{m} x\right) = mm_0 m_1 Q\left(\frac{x}{m}\right),$$

or

$$(1.21) \quad \bar{\bar{Q}}(x) = m^2 Q(m^{-1}x).$$

Therefore from (1.12), (1.20) and (1.21) we have

$$(1.22) \quad Q(x) = m^2 Q(m^{-1}x),$$

which is (1.15) for  $n = 1$ , and  $0 < m < 1$ .

Assume (1.15),  $0 < m < 1$ , is true and from (1.22), with  $m^{-n}x$  on place of  $x$ , we get :

$$(1.23) \quad Q(m^{-(n+1)}x) = m^{-2}Q(m^{-n}x) = m^{-2}(m^{-n})^2 Q(x) = (m^{-(n+1)})^2 Q(x).$$

This formula (1.23), by induction, proves formula (1.15) for  $0 < m < 1$ .

## 2. ULAM STABILITY FOR EULER-LAGRANGE TYPE QUADRATIC FUNCTIONAL EQUATIONS

**Theorem 2.1.** *Let  $X$  and  $Y$  be real normed linear spaces. Assume that  $Y$  is complete. Take  $0 < m = \frac{m_1+m_2}{m_1m_2+1}(m_1a_1^2 + m_2a_2^2) \neq 1$  for any fixed non-zero reals  $a_i$  and positive reals  $m_i$  ( $i = 1, 2$ ). Assume in addition that mappings  $Q : X \rightarrow Y$  and  $f : X \rightarrow Y$  satisfy the Euler-Lagrange type functional equation (1.1) and inequality (1.2), respectively, and conditions*

$$(2.1) \quad \overline{\overline{Q}}(x) = \overline{Q}(x),$$

and

$$(2.2) \quad \left\| \overline{\overline{f}}(x) - \overline{f}(x) \right\| \leq \frac{1}{m_0m_1} \begin{cases} \frac{1}{m}\varepsilon_1, & \text{if } m > 1 \\ m\varepsilon_2, & \text{if } 0 < m < 1, \end{cases}$$

with constants  $\varepsilon_1, \varepsilon_2 \geq 0$  (independent of  $x_1, x_2 \in X$ ), and a positive constant  $m_0 = \frac{m_1m_2+1}{m_1+m_2}$ . Define

$$(2.3) \quad f_n(x) = \begin{cases} m^{-2n}f(m^n x), & \text{if } m > 1 \\ m^{2n}f(m^{-n}x), & \text{if } 0 < m < 1, \end{cases}$$

for all  $x \in X$  and  $n \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ . Then the limit

$$(2.4) \quad Q(x) = \lim_{n \rightarrow \infty} f_n(x)$$

exists for all  $x \in X$  and  $Q : X \rightarrow Y$  is the unique Euler-Lagrange type quadratic mapping, such that

$$(2.5) \quad \begin{aligned} & \|f(x) - Q(x)\| \\ & \leq \frac{\{(m + m_0m_1)(m_1m_2 + 1)|m - 1| + m_0m_2|m^2 - m_1^2|\}c + m(m_1m_2 + 1)|m - 1|\varepsilon}{m_0m_1(m_1m_2 + 1)(m - 1)^2(m + 1)} \\ & = \delta \end{aligned}$$

holds for all  $x \in X$ , with non-negative constants  $c$  and

$$(2.6) \quad \varepsilon = \begin{cases} \varepsilon_1, & \text{if } m > 1 \\ \varepsilon_2, & \text{if } 0 < m < 1 \end{cases} \quad \text{independent of } x \in X.$$

*Proof.* Now claim for  $n \in \mathbb{N}_0$  that inequality

$$(2.7) \quad \|f(x) - f_n(x)\| \leq \begin{cases} \delta_1(1 - m^{-2n}), & \text{if } m > 1 \\ \delta_2(1 - m^{2n}), & \text{if } 0 < m < 1, \end{cases}$$

holds for all  $x \in X$ , where  $\delta_1 = \delta$  for  $m > 1$  and  $\delta_2 = \delta$  for  $0 < m < 1$ :

$$\delta_1 = \frac{\{(m + m_0 m_1)(m_1 m_2 + 1)(m - 1) + m_0 m_2 |m^2 - m_1^2|\} c + m(m_1 m_2 + 1)(m - 1) \varepsilon_1}{m_0 m_1 (m_1 m_2 + 1)(m - 1)^2 (m + 1)} \quad (2.8)$$

$$\delta_2 = \frac{\{(m + m_0 m_1)(m_1 m_2 + 1)(1 - m) + m_0 m_2 |m_1^2 - m^2|\} c + m(m_1 m_2 + 1)(1 - m) \varepsilon_2}{m_0 m_1 (m_1 m_2 + 1)(1 - m)^2 (m + 1)}$$

are two non-negative constants independent of  $x \in X$ . For  $n = 0$ , it is trivial.

Assume  $m > 1$ . From (1.2), with  $x_i = \frac{a_i}{m_0} x$ , ( $i = 1, 2$ ), we obtain

$$\left\| f(mx) + m_1 m_2 f(0) - m m_0 \left[ m_1 f\left(\frac{a_1}{m_0} x\right) + m_2 f\left(\frac{a_2}{m_0} x\right) \right] \right\| \leq c,$$

or

$$(2.9) \quad \left\| \bar{f}(x) - m^{-2} f(mx) - \frac{m_1 m_2}{m^2} f(0) \right\| \leq \frac{c}{m^2},$$

where  $\bar{f}(x) = \frac{m_0}{m} \left[ m_1 f\left(\frac{a_1}{m_0} x\right) + m_2 f\left(\frac{a_2}{m_0} x\right) \right]$ . Besides, from (1.2), with  $x_1 = x, x_2 = 0$ , we get  $\|f(m_1 a_1 x) + m_1 m_2 f(a_2 x) - m m_0 [m_1 f(x) + m_2 f(0)]\| \leq c$ , or

$$(2.10) \quad \left\| f(x) - \bar{\bar{f}}(x) + \frac{m_2}{m_1} f(0) \right\| \leq \frac{c}{m_0 m_1 m},$$

where  $\bar{\bar{f}}(x) = \frac{1}{m_0 m_1 m} [f(m_1 a_1 x) + m_1 m_2 f(a_2 x)]$ . Therefore from (1.4), (2.2), (2.9), (2.10) and triangle inequality we have

$$\begin{aligned} & \left\| f(x) - m^{-2} f(mx) + \frac{m_2(m^2 - m_1^2)}{m_1 m^2} f(0) \right\| \\ & \leq \left\| f(x) - \bar{\bar{f}}(x) + \frac{m_2}{m_1} f(0) \right\| + \left\| \bar{\bar{f}}(x) - \bar{f}(x) \right\| + \left\| \bar{f}(x) - m^{-2} f(mx) - \frac{m_1 m_2}{m^2} f(0) \right\| \\ & \leq \frac{c}{m_0 m_1 m} + \frac{\varepsilon_1}{m_0 m_1 m} + \frac{c}{m^2} = \frac{(m + m_0 m_1) c + m \varepsilon_1}{m_0 m_1 m^2}, \end{aligned}$$

or

$$(2.11) \quad \|f(x) - m^{-2} f(mx)\| \leq \delta_1 (1 - m^{-2}).$$

Assume  $0 < m < 1$ . Similarly, from (1.2), with  $x_i = \frac{a_i}{m_0} \frac{x}{m} = \frac{b_i}{m_0} x$ ,  $b_i = \frac{a_i}{m}$  ( $i = 1, 2$ ), we obtain  $\|f(x) + m_1 m_2 f(0) - m m_0 \left[ m_1 f\left(\frac{b_1}{m_0} x\right) + m_2 f\left(\frac{b_2}{m_0} x\right) \right]\| \leq c$ , or

$$(2.12) \quad \|f(x) - \bar{f}(x) + m_1 m_2 f(0)\| \leq c,$$

where  $\bar{f}(x) = m_0 m \left[ m_1 f\left(\frac{b_1}{m_0} x\right) + m_2 f\left(\frac{b_2}{m_0} x\right) \right]$ . Besides, from (1.2), with  $x_1 = \frac{x}{m}, x_2 = 0$ , we get  $\|f(m_1 b_1 x) + m_1 m_2 f(b_2 x) - m m_0 \left[ m_1 f\left(\frac{x}{m}\right) + m_2 f(0) \right]\| \leq c$ , or

$$(2.13) \quad \left\| \bar{\bar{f}}(x) - m^2 f\left(m^{-1} x\right) - \frac{m_2}{m_1} m^2 f(0) \right\| \leq \frac{m}{m_0 m_1} c,$$

where  $\bar{\bar{f}}(x) = \frac{m}{m_0 m_1} [f(m_1 b_1 x) + m_1 m_2 f(b_2 x)]$ . Therefore from (1.4), (2.2), (2.12), (2.13) and triangle inequality we have

$$\begin{aligned} & \left\| f(x) - m^2 f(m^{-1}x) + \frac{m_2(m_1^2 - m^2)}{m_1} f(0) \right\| \\ & \leq \|f(x) - \bar{f}(x) + m_1 m_2 f(0)\| + \|\bar{f}(x) - \bar{\bar{f}}(x)\| + \left\| \bar{\bar{f}}(x) - m^2 f(m^{-1}x) - \frac{m_2}{m_1} m^2 f(0) \right\| \\ & \leq \frac{(m + m_0 m_1)c + m\varepsilon_2}{m_0 m_1}, \end{aligned}$$

or

$$(2.14) \quad \|f(x) - m^2 f(m^{-1}x)\| \leq \delta_2 (1 - m^2).$$

Assume  $m > 1$ . From (2.11), with  $m^i x$  ( $i = 1, 2, \dots, n$ ), on place of  $x$ , and the triangle inequality, we have, *without induction*

$$\begin{aligned} & \|f(x) - f_n(x)\| = \|f(x) - m^{-2n} f(m^n x)\| \\ & \leq \|f(x) - m^{-2} f(mx)\| + m^{-2} \|f(mx) - m^{-2} f(m^2 x)\| \\ & + \dots + m^{-2(n-1)} \|f(m^{n-1}x) - m^{-2} f(m^n x)\| \\ & \leq \delta_1 (1 + m^{-2} + \dots + m^{-2(n-1)}) (1 - m^{-2}) = \delta_1 (1 - m^{-2n}). \end{aligned}$$

Similarly if we assume  $0 < m < 1$ , we have from (2.14) that

$$\begin{aligned} & \|f(x) - f_n(x)\| = \|f(x) - m^{2n} f(m^{-n}x)\| \\ & \leq \delta_2 (1 + m^2 + \dots + m^{2(n-1)}) (1 - m^2) = \delta_2 (1 - m^{2n}). \end{aligned}$$

Therefore we prove inequality (2.7).

Claim now that the sequence  $\{f_n(x)\}$  converges. To do this it suffices to prove that it is a Cauchy sequence. Inequality (2.7) is involved.

In fact, if  $m > 1$  and  $i > j > 0$ , and  $h_1 = m^j x$ , we have

$$\begin{aligned} & \|f_i(x) - f_j(x)\| \\ & = m^{-2j} \|m^{-2(i-j)} f(m^{i-j} h_1) - f(h_1)\| \\ & \leq \delta_1 m^{-2j} (1 - m^{-2(i-j)}) \\ & = \delta_1 (m^{-2j} - m^{-2i}) \\ & < \delta_1 m^{-2j} \xrightarrow{j \rightarrow \infty} 0. \end{aligned}$$

Similarly, if  $0 < m < 1$ , and  $h_2 = m^{-j} x$ , we have :

$$\|f_i(x) - f_j(x)\| = m^{2j} \|m^{2(i-j)} f(m^{-(i-j)} h_2) - f(h_2)\| < \delta_2 m^{2j} \xrightarrow{j \rightarrow \infty} 0.$$

Thus we can *define* a mapping  $Q : X \rightarrow Y$ , by (2.4).

Claim that from (1.2) and (2.4) we can get (1.1), or equivalently that the afore-mentioned well-defined mapping  $Q : X \rightarrow Y$  is *Euler-Lagrange type quadratic*. In fact, it is clear from the functional inequality (1.2) and the limit (2.4) with  $m > 1$  that

$$\begin{aligned} & m^{-2n} \|f(m_1 a_1 m^n x_1 + m_2 a_2 m^n x_2) + m_1 m_2 f(a_2 m^n x_1 - a_1 m^n x_2) \\ & - (m_1 a_1^2 + m_2 a_2^2) [m_1 f(m^n x_1) + m_2 f(m^n x_2)]\| \leq m^{-2n} c, \end{aligned}$$

or

$$\begin{aligned} & \|f_n(m_1a_1x_1 + m_2a_2x_2) + m_1m_2f_n(a_2x_1 - a_1x_2) \\ & - (m_1a_1^2 + m_2a_2^2) [m_1f_n(x_1) + m_2f_n(x_2)]\| \\ & \leq m^{-2n}c \xrightarrow[n \rightarrow \infty]{} 0, \end{aligned}$$

or

$$\begin{aligned} & \|Q(m_1a_1x_1 + m_2a_2x_2) + m_1m_2Q(a_2x_1 - a_1x_2) - (m_1a_1^2 + m_2a_2^2) [m_1Q(x_1) + m_2Q(x_2)]\| \\ & = 0 \end{aligned}$$

or the mapping  $Q$  satisfies (1.1) if  $m > 1$ . Similarly, from (1.2) and (2.4) we get that  $Q$  satisfies (1.1) if  $0 < m < 1$ . Therefore  $Q$  satisfies (1.1) if  $0 < m \neq 1$ , completing the proof that  $Q$  is *Euler-Lagrange type quadratic mapping* in  $X$ .

It is now clear from inequality (2.7) with  $n \rightarrow \infty$ , as well as formula (2.4) that the required inequality (2.5) holds in  $X$ . This completes ***the existence proof*** of the above-mentioned Theorem 2.1.

We claim that  $Q$  is ***unique***. Let  $Q' : X \rightarrow Y$  be another Euler-Lagrange type quadratic mapping satisfying (2.5). Then  $Q' = Q$ .

In fact, assume  $m > 1$ . Remember both  $Q$  and  $Q'$  satisfy (1.15). Then for every  $x \in X$  and  $n \in \mathbb{N}_0$ ,

$$\begin{aligned} & \|Q(x) - Q'(x)\| \\ & \leq m^{-2n} \{ \|Q(m^n x) - f(m^n x)\| + \|Q'(m^n x) - f(m^n x)\| \} \\ & \leq 2\delta m^{-2n} \xrightarrow[n \rightarrow \infty]{} 0, \end{aligned}$$

or  $Q(x) = Q'(x)$ . Similarly we establish uniqueness results if  $0 < m < 1$ . This completes the proof of *the uniqueness* and ***the Ulam stability*** for Euler-Lagrange type quadratic functional equations of the form (1.1). ■

**Corollary 2.2.** *Let  $X$  and  $Y$  be real normed linear spaces. Assume that  $Y$  is complete. Take  $m_1 = m_2 = 1 : 0 < m = a_1^2 + a_2^2 \neq 1$  and  $m_0 = 1$  for any fixed non-zero reals  $a_i$  ( $i = 1, 2$ ). Define functions  $f_n = f_n(x)$  as in (2.3). Then the limit in (2.4) exists for all  $x \in X$  and  $Q : X \rightarrow Y$  is the unique Euler-Lagrange type quadratic mapping such that*

$$(2.15) \quad \|f(x) - Q(x)\| \leq \frac{3}{2} \frac{1}{|m-1|} c (= \delta)$$

for all  $x \in X$  with constant  $c \geq 0$  (independent of  $x \in X$ ).

Note that in this case there is ***no constant***  $\varepsilon$  in the right-hand side of (2.15) because  $\overline{\overline{f}}(x) = \overline{f}(x)$ . Besides  $\delta$  given by (2.15) is sharper than the corresponding one in [21, 22] which is of the form

$$\begin{aligned} \delta &= \frac{1}{2} \frac{c}{(m-1)^2(m+1)} \begin{cases} 3m^2 - 1, & \text{if } m > 1 \\ 3 - m^2, & \text{if } 0 < m < 1 \end{cases} \\ &\geq \frac{1}{2} \frac{c}{(m-1)^2(m+1)} \begin{cases} 3m^2 - 3, & \text{if } m > 1 \\ 3 - 3m^2, & \text{if } 0 < m < 1 \end{cases} \\ &= \frac{3}{2} \frac{1}{|m-1|} c. \end{aligned}$$



If  $a_1 = a_2 = 1$ , then  $m = 2$  and from (2.15) we have  $\delta = \frac{3c}{2}$ . We note that in this case **a sharper constant**  $\delta = \frac{c}{2}$  may be found, if new substitution  $x_1 = x_2 = x$  is applied in (1.2), because  $\overline{\overline{f}}(x) = \overline{f}(x) = f(x)$  [21, 22].

## REFERENCES

- [1] J. ACZEL, *Lectures on Functional Equations and their Applications*, Academic Press, New York and London, 1966.
- [2] C. BORELLI and G.L. FORTI, On a general Hyers-Ulam stability result, *Internat. J. Math. Sci.*, **18** (1995), 229-236.
- [3] D.G. BOURGIN, Classes of transformations and bordering transformations, *Bull. Amer. Math. Soc.*, **57** (1951), 223-237.
- [4] P.W. CHOLEVA, Remarks on the stability of functional equations, *Aequationes Math.*, **27** (1984), 76- 86.
- [5] ST. CZERWIK, On the stability of the quadratic mapping in normed spaces, *Abh. Math. Sem. Univ Hamburg*, **62** (1992), 59-64.
- [6] H. DRLJEVIC, On the stability of the functional quadratic on A-orthogonal vectors, *Publ. Inst. Math. (Beograd) (N.S.)*, **36**(50) (1984), 111-118.
- [7] I. FENYO, Osservazioni su alcuni teoremi di D.H. Hyers, *Istit. Lombardo Accad. Sci. Lett. Rend.*, **A 114** (1980), (1982) , 235-242.
- [8] I. FENYO, On an inequality of P.W.Cholewa. In: *General Inequalities*, **5**. [Internat. Schriftenreihe Numer. Math., Vol. 80]. Birkhäuser, Basel-Boston, MA, 1987, pp. 277-280.
- [9] G.L. FORTI, Hyers-Ulam stability of functional equations in several variables, *Aequationes Math.*, **50** (1995), 143-190.
- [10] Z. GAJDA and R. GER, Subadditive multifunctions and Hyers-Ulam stability. In: *General Inequalities*, **5**. [Internat. Schriftenreihe Numer. Math., Vol. 80]. Birkhäuser, MA, 1987.
- [11] P. GAVRUTA, An answer to a question of John M. Rassias concerning the stability of Cauchy equation. In: *Advances in Equations and Inequalities*, Hadronic Math. Series, U.S.A., 1999, 67-71.
- [12] P.M. GRUBER, Stability of isometries, *Trans. Amer. Math. Soc.*, U.S.A., **245** (1978), 263-277.
- [13] D.H. HYERS, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci.* **27** (1941), 222-24: The stability of homomorphisms and related topics, "Global Analysis-Analysis on Manifolds", *Teubner - Texte zur Mathematik*, **57** (1983), 140-153.
- [14] S.-M. JUNG, On the Hyers-Ulam stability of the functional equations that have the quadratic property, *J. Math. Anal. & Appl.*, **222** (1998), 126-137.
- [15] PL. KANNAPPAN, Quadratic functional equation and inner product spaces, *Results Math.*, **27** (1995), 368-372.
- [16] J.M. RASSIAS, On approximation of approximately linear mappings by linear mappings, *J. Funct. Anal.* **46** (1982), 126-130.
- [17] J.M. RASSIAS, On approximation of approximately linear mappings by linear mappings, *Bull. Sc. Math.* **108** (1984), 445-446.
- [18] J.M. RASSIAS, Solution of a problem of Ulam, *J. Approx. Th.* **57** (1989), 268-273.
- [19] J.M. RASSIAS, On the stability of the Euler-Lagrange functional equation, *Chin. J. Math.* **20** (1992), 185-190.

- [20] J.M. RASSIAS, On the stability of a multi-dimensional Cauchy type functional equation, "Geometry, Analysis and Mechanics", World Sci. Publ. Co., 1994, 365-376.
- [21] J.M. RASSIAS, On the stability of the general Euler-Lagrange functional equation, *Demonstr. Math.* **29** (1996), 755-766.
- [22] J.M. RASSIAS, Solution of the Ulam stability problem for Euler-Lagrange quadratic mappings, *J. Math. Anal. Applications* **220** (1998), 613-639.
- [23] J.M. RASSIAS, Generalization of the Euler theorem to heptagons leading to a quadratic vector identity, *Advances in Equations and Inequalities*, Hadronic Press (2000), 179-183.
- [24] J.M. RASSIAS, Solution of the Ulam stability problem for an Euler type quadratic functional equation, *Southeast Asian Bull. Math.*, Springer-Verlag, **26**(2002), 101-112.
- [25] J.M. RASSIAS, On the Ulam stability of mixed type mappings on restricted domains, *J. Math. Anal. Appl.* **276**(2002), 747-762.
- [26] J.M. RASSIAS and M.J. RASSIAS, On the Ulam stability of Jensen and Jensen type mappings on restricted domains, *J. Math. Anal. Appl.* **281**(2003) 516-524.
- [27] J.M. RASSIAS, Asymptotic behavior of mixed type functional equations, *Austral. J. Math. Anal. Applications*, **1**(2004), Issue 1, 1-21
- [28] S.M. ULAM, "A Collection of Mathematical Problems", Interscience Publishers, Inc., New York, 1968, p. 63.